



York Teaching Hospital
NHS Foundation Trust

Information for Patients with Diabetes having
Lower Endoscopic Investigations requiring full
bowel preparation

Afternoon Appointment

Information for patients, relatives and carers

① For more information, please contact:

Diabetes and Endocrine Unit

The York Hospital
Wigginton road, York, YO31 8HE

Tel: 01904 726510

Scarborough Hospital
Woodlands Drive, Scarborough, YO12 6QL

Tel: 01723 342274

Caring with pride

Contents	Page
General advice	3
What to do with your medication.....	4
Monitoring your blood glucose levels.....	4
What is a hypo?.....	5
What to do if you have a hypo	5
What to do after your procedure	6
What type of insulin do I take?.....	7
Insulin recommendations for Afternoon Appointment.....	8
Day before the procedure:	8
Day of the procedure	9
Non – insulin medication recommendation for afternoon appointment	10
Day before the procedure	10
Day of the procedure	10
Tell us what you think of this leaflet	11
Teaching, Training and Research	11
Patient Advice and Liaison Service (PALS).....	11

General advice

For a successful examination to be carried out, it is very important that the bowel is completely clean and you follow the guidance below. If the bowel is not completely clean and /or the guidance is not followed correctly the examination may have to be repeated.

Some investigations required you to fast for extended periods of time to ensure the bowel is completely clean. For people with diabetes, any procedure that causes you to miss a meal or change your usual meal plan will require special planning to safely manage your blood glucose and your medications may need some adjustment.

If you normally take insulin for your diabetes you may need to replace solid food with sugary drinks, to prevent your blood glucose level dropping too low. Examples for meal replacement could include:

- 300 millimetres (mls) full sugar fizzy drink
- 300 - 400mls clear apple juice

What to do with your medication.

For people who take insulin injections, your usual doses may need some adjustment on the day of your bowel preparation and the day of the procedure. The charts on pages 8-10 will help you manage your diabetes treatment if taking insulin when undergoing your bowel preparation or fasting period.

For people with Type 2 diabetes on tablets or injectable treatment see guidance on page 10.

Monitoring your blood glucose levels

It is advisable to test your blood glucose regularly **if you normally test**. You may need to test more often than normal. We would advise testing before each meal, before bed, and if you feel at any time that your blood glucose level is getting low. If you don't normally test your blood glucose levels then you don't need to test during this period.

When coming to the hospital please bring your medication and any equipment you need, as well as a light meal (for example a sandwich). Once you have recovered from the procedure, you will be able to eat, drink and resume your normal diabetes medications.

What is a hypo?

A hypo or low blood glucose level is where your blood glucose level falls below the normal range – usually less than 4mmol/L. This is often associated with symptoms such as feeling sweaty, shaky, dizzy and heart racing. You might be at risk of a hypo if you take insulin or tablets such as gliclazide, glipizide or glimepiride (metformin on its own does not cause hypos).

What to do if you have a hypo

You need to treat your hypo immediately by taking rapid acting carbohydrate for example:

- 150mls – full sugar fizzy drink
- Five to seven Dextrose tablets
- Four to five Glucotabs
- 60ml bottle Glucojuice
- 200mls clear apple juice.

Recheck your blood glucose level 10 - 15 minutes later and repeat with more rapid acting carbohydrate if your result is still less than 4mmol/L.

Once your blood glucose level has risen above 4mmol/L recheck hourly to ensure your blood glucose does not drop too low again.

What to do after your procedure

The nursing staff will inform you when it is safe to eat and drink after your procedure. You can resume your normal diabetes medications once you have recovered from the procedure and are eating and drinking as normal.

When you get home your blood glucose levels may vary for a short time, but this should settle quickly. If, however, you take the diabetes medication - Metformin, Dapagliflozin, Canagliflozin or Empagliflozin and become unwell following discharge home and are vomiting, **stop** these medications and consult your GP or diabetes care provider for advice.

These are general guidelines, if you have any concerns please call the diabetes team for further advice.

What type of insulin do I take?

To help you work out what to do with your insulin, you need to know what type of insulin you take. Here is a list to help you.

Fast acting insulin	Mixed insulin	Long acting insulin
Humulin S Actrapid Novorapid Humalog Apidra Fiasp	Humulin M3 Humalog Mix 25 Humalog Mix 50 Novomix 30 Insuman Comb 15 Insuman Comb 25 Insuman Comb 50	Humulin I Insulatard Insuman basal Lantus Abasaglar Levemir Toujeo Tresiba100 Tresiba 200

Insulin recommendations for Afternoon Appointment

Day before the procedure

Follow dietary and bowel preparation as instructed

Drink additional clear fluids (e.g. black tea or coffee)
some of it sugary (e.g. clear fruit juice) or clear jelly
puddings (not red jelly)

Test your blood glucose **at least four times a day**

Fast acting or mixed insulin	Long acting insulin
Take half usual dose	Take as normal if you have Type 1 diabetes. Reduce the dose/s by 20% (one fifth) if you have Type 2 diabetes.

Insulin recommendations for Afternoon Appointment

Day of the procedure

Continue bowel preparation as instructed

Drink additional clear fluids (e.g. black tea or coffee) some of it sugary (e.g. clear fruit juice) or clear jelly puddings (not red jelly), until two hours before your appointment

Test your blood glucose **every two hours**

Fast acting insulin	Mixed insulin	Long acting insulin
Omit at breakfast and lunch	Take half breakfast dose and omit at lunchtime if taken at this time	Take as normal if you have Type 1 diabetes Reduce the morning dose, if taken at this time, by 20% (one fifth) if you have Type 2 diabetes

Non – insulin medication recommendation for afternoon appointment

Day before the procedure

Omit all diabetes tablets or injectable diabetes treatment
(Follow separate guidance for insulin)

Day of the procedure

Take these medications as normal	Omit all doses and resume when eating and drinking normally	Take as normal unless taken at lunchtime in which case omit lunchtime dose
Pioglitazone Byetta (Exenatide) Trulicity(Dulaglutide) Victosa (liraglutide) Lyxumia(Lixisenatide) Bydureon Linagliptin Saxagliptin	Gliclazide Glipizide Glimepiride Dapagliflozin Empagliflozin Canagliflozin Acarbose Repaglinide Neteglinide	Metformin

Tell us what you think of this leaflet

We hope that you found this leaflet helpful. If you would like to tell us what you think, please contact:

Tara Kadis, Diabetes & Endocrine Unit,
The York Hospital, telephone 01904 726510 or email
tara.kadis@york.nhs.uk.

Teaching, Training and Research

Our Trust is committed to teaching, training and research to support the development of health and healthcare in our community. Healthcare students may observe consultations for this purpose. You can opt out if you do not want students to observe. We may also ask you if you would like to be involved in our research.

Patient Advice and Liaison Service (PALS)

PALS offers impartial advice and assistance to patients, their relatives, friends and carers. We can listen to feedback (positive or negative), answer questions and help resolve any concerns about Trust services.

PALS can be contacted on 01904 726262, or email
pals@york.nhs.uk.

An answer phone is available out of hours.

Please telephone or email if you
require this information in a different
language or format

如果你要求本資訊是以不同的語言
或版式提供，請致電或寫電郵

Jeżeli niniejsze informacje potrzebne
są w innym języku lub formie,
należy zadzwonić lub wysłać
wiadomość e-mail

Bu bilgileri değişik bir lisanda ya da
formatta istiyorsanız lütfen telefon
ediniz ya da e-posta gönderiniz



01904 725566

email: access@york.nhs.uk



Braille



Audio e.g.
CD



Large print



Electronic

Owner	Tara Kadis – Diabetes Team
Date first issued	October 2015
Review Date	May 2020
Version	2 (issued June 2018)
Approved by	Diabetes and Endocrine Team
Document Reference	PIL 972 v2

© 2018 York Teaching Hospital NHS Foundation Trust. All Rights reserved