

## Tell us what you think

We hope that you found this leaflet helpful. If you would like to tell us what you think, please contact Mr R H Taylor, Consultant Ophthalmologist, The York Hospital, Wigginton Road, York, YO31 8HE or telephone 01904 725612.

## Teaching, training & research

Our Trust is committed to teaching, training and research to support the development of health and healthcare in our community. Healthcare students may observe consultations for this purpose. You can opt out if you do not want students to observe. We may also ask you if you would like to be involved in our research.

## Patient Advice and Liaison Service (PALS)

PALS offers impartial advice and assistance to patients, their relatives, friends and carers. We can listen to feedback (positive or negative), answer questions and help resolve any concerns about Trust services. PALS can be contacted on 01904 726262, or email [pals@york.nhs.uk](mailto:pals@york.nhs.uk). An answer phone is available out of hours.

Providing care together in York,  
Scarborough, Bridlington, Malton,  
Selby and Easingwold communities



York Teaching Hospital  
NHS Foundation Trust

Please telephone or email if you require this information in a different language or format

如果你要求本資訊是以不同的語言或版式提供，請致電或寫電郵

Jeżeli niniejsze informacje potrzebne są w innym języku lub formie, należy zadzwonić lub wysłać wiadomość e-mail

Bu bilgileri değişik bir lisanda ya da formatta istiyorsanız lütfen telefon ediniz ya da e-posta gönderiniz



**01904 725566**

email: [access@york.nhs.uk](mailto:access@york.nhs.uk)



Braille



Audio e.g.  
CD



Large print



Electronic

Owner

Mr R H Taylor,  
Consultant Ophthalmologist

Date first issued

February 2003

Review Date

September 2019

Version

4 (reissued October 2017)

Approved by

Ophthalmology Team

Document Reference

PIL 160 v4.2

© 2017 York Teaching Hospital NHS Foundation Trust

All rights reserved

# Strabismus (Squint) Surgery

Information for patients,  
relatives and carers

① For more information contact:

Mr Taylor's Secretary  
Tel: 01904 725612

**Ophthalmology Department**

The York Hospital  
Wigginton Road, York, YO31 8HE

Caring with pride

## Strabismus (Squint) Surgery

The intended benefits and possible complications relating to strabismus surgery are listed in this leaflet. No list is complete, as with any surgery there are unusual risks that are too numerous to list. The final result cannot be guaranteed. If you have any questions after reading this leaflet please discuss these with your doctor.

### What are the intended benefits of strabismus surgery?

- To reduce/eliminate double vision
- To bring areas of single vision into the centre
- To straighten the eyes
- To reduce asthenopic symptoms (for example headache, eye pain)
- To reduce or eliminate compensatory head posture
- To promote binocularity
- To reduce visual confusion (loss of suppression)
- To reduce requirement for prism, patch or occlusion
- To improve visual field of vision

### Are there any possible complications during the operation?

(These may lead to loss of sight, loss of the eye or a requirement for further surgery)

- Incorrect eye or muscle operated on
- Damage to the eye or vision
- Damage to the eye with suture needle or other sharp object
- Bleeding inside or outside the eye
- Loss of a muscle or muscle torn in two

### What are the possible complications after my operation?

(These may lead to loss of sight, loss of the eye or a requirement for further surgery)

- Swelling of the eye or eyelids
- Redness of the 'white' of the eye that persists beyond two months
- Prominent scarring of the 'white' of the eye
- Under or over correction of the eye position leading in some cases to further squint surgery (risk 10% in average case)
- Deterioration of double vision, or new double vision, which is disabling
- Infection outside or inside the eye
- Allergy to any medication used

**In an average case, the likelihood of a serious complication (permanent reduced vision or troublesome double vision) is one in 2,400.**