

Afternoon Appointment Insulin recommendations

Day before Procedure

If you take long acting insulin such as:

Lantus, Levemir, Humulin I, Insuman Basal, Abasaglar, Toujeo, Tresiba

Reduce your dose of insulin by 20% the day before your procedure

Day of Procedure

Type of Insulin	What to do
<p>Once daily (long acting insulin) evening injection</p> <p>Lantus Levemir Humulin I Insuman Basal Abasaglar Toujeo Tresiba 100 or 200</p>	<p>No dose adjustment necessary</p> <p>You will be able to have usual evening insulin after the procedure</p>
<p>Once daily (long acting insulin) morning injection</p> <p>Lantus Levemir Humulin I Insuman Basal Abasaglar Toujeo Tresiba 100 or 200</p>	<p>Reduce morning dose by 20%</p>
<p>Twice daily insulin</p> <p>Humulin M3 Novomix 30 Humalog Mix 25 Humalog Mix 50 Levemir or Lantus given twice a day</p>	<p>Half your usual morning dose.</p> <p>You can take usual evening dose after the procedure as normal</p>
<p>Multiple daily injection regime This includes an injection of short acting insulin with meals and once or twice daily background insulin</p>	<p>Take usual morning dose of short acting insulin if eating breakfast. Omit lunchtime dose</p> <p>Reduce morning long acting insulin dose if taken in morning by 20%</p>

**Afternoon appointment
Non – insulin medication
recommendations**

Day before procedure

Omit the SGLT2 inhibitor (e.g. dapagliflozin, canagliflozin, empagliflozin, ertugliflozin)
Take all other medication as normal the day before

Day of procedure

Medication	What to do
Metformin	Take as normal in morning Omit lunchtime dose if taken Take evening dose as usual
Sulfonylureas (e.g. gliclazide, glipizide, glimepiride)	Omit all doses taken on day of procedure
Meglitinides (e.g. repaglinide or nateglinide)	Give morning dose if eating breakfast
Pioglitazone	Take as normal
DPP4 inhibitors (e.g. sitagliptin, linagliptin, alogliptin, saxagliptin)	Take as normal
SGLT2 inhibitors (e.g. dapagliflozin, canagliflozin, empagliflozin, ertugliflozin)	Omit dose
GLP1 therapy (e.g. Byetta, Victoza, trulicity, Ozempic, Rybelsus, Lyxumia, MOUNJARO)	Take as normal

Remember to test blood glucose more regularly if you usually test,
and that your
glucose level may be a little erratic for a few days after the procedure.

Keep hypo treatment with you in case of low blood glucose levels
(glucose level under 4 mmol/L)

For further advice contact the Diabetes Team:
York (01904) 725789
Scarborough (01723) 342274

Tell us what you think of this leaflet

We hope that you found this leaflet helpful. If you would like to tell us what you think, please contact Nic Lloyd-Jones, Diabetes and Endocrine Unit, York Hospital, telephone 01904 725789 or email Nicola.lloyd-jones1@nhs.net

Patient Advice and Liaison Service (PALS)

PALS offers impartial advice and assistance to patients, their relatives, friends and carers. We can listen to feedback (positive or negative), answer questions and help resolve any concerns about Trust services.

PALS can be contacted on 01904 726262, or email yhs-tr.patientexperienceteam@nhs.net

An answer phone is available out of hours.

Teaching, training and research

Our Trust is committed to teaching, training and research to support the development of health and healthcare in our community. Healthcare students may observe consultations for this purpose. You can opt out if you do not want students to observe. We may also ask you if you would like to be involved in our research.

Leaflets in alternative languages or formats

If you would like this information in a different format, including braille or easy read, or translated into a different language, please speak to a member of staff in the ward or department providing your care.

Patient Information Leaflets can be accessed via the Trust's Patient Information Leaflet website: www.yorkhospitals.nhs.uk/your-visit/patient-information-leaflets/

Owner	Nic Lloyd-Jones – Diabetes Team
Date first issued	October 2018
Review Date	January 2029
Version	3 (issued January 2026)
Approved by	Diabetes and Endocrine Team
Document Reference	PIL 1268 v3
© 2026 York and Scarborough Teaching Hospitals NHS Foundation Trust. All Rights reserved	

www.yorkhospitals.nhs.uk