# Diabetes Medication Guidelines for people with diabetes requiring Upper GI Endoscopy



# Morning Appointment Insulin recommendations

#### **Day before Procedure**

#### If you take long acting insulin such as:

Lantus, Levemir, Humulin I, Insuman Basal, Abasaglar, Toujeo, Tresiba Reduce your dose of insulin by 20% the day before your procedure

#### **Day of Procedure**

| Type of Insulin                     | What to do                           |
|-------------------------------------|--------------------------------------|
| Once daily (long acting insulin)    | Do not need to do anything – will be |
| evening injection                   | able to have usual evening insulin   |
| Lantus                              | after the procedure                  |
| Levemir                             |                                      |
| Humulin I                           |                                      |
| Insuman Basal                       |                                      |
| Abasaglar                           |                                      |
| Toujeo                              |                                      |
| Tresiba 100 or 200                  |                                      |
| Once daily (long acting insulin )   |                                      |
| morning injection                   |                                      |
| Lantus                              | Reduce dose by 20%                   |
| Levemir                             |                                      |
| Humulin I                           |                                      |
| Insuman Basal Abasaglar             |                                      |
| Toujeo                              |                                      |
| Tresiba 100 or 200                  |                                      |
| Twice daily insulin                 |                                      |
| Humulin M3                          | Half usual morning dose.             |
| Novomix 30                          |                                      |
| Humalog Mix 25                      | You can take usual evening dose      |
| Humalog Mix 50                      | after the procedure as normal        |
| Levemir or Lantus given twice a day |                                      |
| Three times daily mixed insulin     |                                      |
| Humulin M3                          | Half usual morning dose              |
| Humalog Mix 25                      | Omit lunch time dose                 |
| Humalog Mix 50                      | Resume normal if eating a normal     |
| NovoMix 30                          | meal. If small give half usual dose  |
| THO VOIVILA GO                      |                                      |
|                                     |                                      |

# Multiple daily injection regime This includes an injection of short acting insulin with meals and once or twice daily background insulin Reduce morning long acting insulin dose if taken in morning by 20%

Diabetes Medication Guidelines for people with diabetes requiring Upper GI Endoscopy



Morning appointment
Non – insulin medication
recommendations

#### Day before the procedure

Omit the SGLT2 inhibitor (e.g dapagliflozin, canagliflozin, ertugliflozin)

Take all other medication as normal the day before

#### Day of the procedure

| Medication  | What to do   |
|---|--|
| Metformin   | Take as normal in morning - if a dose is taken at lunchtime – omit this dose             |
| Sulfonylureas<br>(e.g. gliclazide, glipizide, glimepiride)                      | Omit the morning dose.  Take usual evening dose if taken twice a day following procedure |
| Meglitinides (e.g. repaglinide or nateglinide)                                  | Omit morning dose  |
|   | Take usual evening dose if taken twice a day following procedure                         |
| Pioglitazone  | Take as normal   |
| <b>DPP4 inhibitors</b> (e.g. sitagliptin, linagliptin, alogliptin, saxagliptin) | Take as normal   |
| SGLT2 inhibitors<br>(e.g. dapagliflozin, canagliflozin,<br>empagliflozin)       | Omit dose  |
| GLP1 injections (e.g. exenatide, liraglutide lixisenatide, dulaglutide)         | Take as normal   |

Remember to test blood glucose more regularly if you usually test, and that your glucose level may be a little erratic for a few days after the procedure.

Keep hypo treatment with you in case of low blood glucose levels (glucose level under 4 mmol/L)

For further advice contact the Diabetes Team: York (01904) 726510 Scarborough (01723) 342274

# Tell us what you think of this leaflet

We hope that you found this leaflet helpful. If you would like to tell us what you think, please contact Tara Kadis, Diabetes and Endocrine Unit, The York Hospital, telephone 01904 726510 or email tara.kadis@york.nhs.uk.

### Leaflets in alternative languages or formats

Please telephone or email if you require this information in a different language or format, for example Braille, large print or audio.

#### 如果你要求本資 不同的 或 式提供, 電或發電

Jeżeli niniejsze informacje potrzebne są w innym języku lub formacje, należy zadzwonić lub wysłać wiadomość e-mail

Bu bilgileri değişik bir lisanda ya da formatta istiyorsanız lütfen telefon ediniz ya da e-posta gönderiniz

Telephone: 01904 725566 Email: access@york.nhs.uk

## Patient Advice and Liaison Service (PALS)

PALS offers impartial advice and assistance to patients, their relatives, friends and carers. We can listen to feedback (positive or negative), answer questions and help resolve any concerns about Trust services. PALS can be contacted on 01904 726262, or email pals@york.nhs.uk. An answer phone is available out of hours.

Owner Tara Kadis – Diabetes Team

Date first issued October 2018 Review Date June 2025

Version 2 (issued June 2022)

Approved by Diabetes and Endocrine Team

Document Reference PIL 1269 v2

© 2022 York and Scarborough Teaching Hospitals NHS Foundation Trust. All Rights reserved

www.yorkhospitals.nhs.uk