

Orthopaedic Department York teaching hospital

This is a follow-up letter to your recent telephone consultation with the fracture care team, explaining the ongoing management of your injury. Your case has been reviewed by an Orthopaedic Consultant (Bone Specialist) and Fracture Care Physiotherapist.

You have sustained a fracture to one of the bones in your elbow (Proximal radius).

Healing: This normally takes approximately 6 weeks to heal.

Pain: Your elbow may be swollen. Take pain killers as prescribed.

Using your arm: It is important to keep the elbow moving to prevent stiffness.

However you should avoid any heavy lifting for the full 6 weeks.

X-Ray: You will need to have a repeat x-ray at approximately 2-3 weeks post injury

to check there has been no movement of the fracture.

Please can you contact us using the e-mail or telephone number at the top of this letter to tell us when this has been done. We will then assess your x-ray

the next day.

Area of Injury:



If you are worried that you are unable to follow this rehabilitation plan, or have any questions, then please phone the Fracture Care Team for advice.

Or if you are experiencing pain and symptoms anywhere else, other than at the site of the original injury or surrounding area, please also get in touch using the telephone or e-mail details at the top of this letter.

Email: virtualfractclinic@york.nhs.uk



Please follow the management plan below.

What to expect:

Weeks since injury	Rehabilitation plan
0-2	 ✓ Wear sling for comfort only. ✓ Take your arm out of the sling and practise moving the elbow (see below). ✓ Do the exercise below every hour in the daytime. You can start straight away.
2-6	 ✓ Try not to use the sling and do normal light activities with the arm. ✗ Don't lifting more than the weight of a cup of tea for six weeks. ✓ Continue with the exercises.
6 -12	 The fracture will be healed Resume normal, day-to-day, activities but be guided by any pain you get. Heavy tasks may cause discomfort.

Advice for a new injury:

Cold packs: A cold pack (ice pack or frozen peas wrapped in a damp towel) can provide short-

term pain relief. Apply this to the sore area for up to 15 minutes, every few hours.

The ice must never be in direct contact with the skin.

Rest: Try to rest your arm for the first 24-72 hours. However, it is important to maintain

movement in your elbow. Gently move your elbow following the exercises shown. These should not cause too much pain. This will ensure your elbow does not become

stiff and it will help the healing process.

Smoking advice:

Medical evidence suggests that smoking prolongs fracture healing time. In extreme cases it can stop healing altogether. It is important that you consider this information with relation to your recent injury. Stopping smoking during the healing phase of your fracture will help ensure optimal recovery from this injury.

For advice on smoking cessation and local support available, please refer to the following website: http://smokefree.nhs.uk or discuss this with your GP.

York Virtual Fracture Clinic Contact details:

Phone: 01904 726575

Email: virtualfractclinic@york.nhs.uk




Exercises

Initial Exercises to be done 5-6 times a day













Elbow Bend to Straighten

Finger and wrist flexion and extension

Open and close your hand as shown 10-15 times. Then move your wrist up and down 10-15 times.

> Bend and straighten your elbow as far as you can without pain. You should not feel more than a mild to moderate stretch. You can use your other arm to assist if necessary.

Repeat 10 - 15 times if there is no increase in pain.





Forearm Rotations

Begin this exercise with your elbow at your side and bent. Slowly turn your palm up and down as far as you can go without pain. You should not feel more than a mild to moderate stretch. You can use your other arm to assist if necessary.

Repeat 10 - 15 times if there is no increase in pain.



Phone: 01904 726575 **Email**: virtualfractclinic@york.nhs.uk